

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 January 2002 (31.01.2002)

PCT

(10) International Publication Number
WO 02/08172 A1

- (51) International Patent Classification⁷: C07C 231/02, 247/12 (74) Agent: R.A. EGLI & CO.; Horneggstrasse 4, Postfach, CH-8034 Zurich (CH).
- (21) International Application Number: PCT/CH01/00400 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 26 June 2001 (26.06.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 1464/00 25 July 2000 (25.07.2000) CH (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*):
SPEEDEL PHARMA AG [CH/CH]; Hirschgässlein 11, CH-4051 Basel (CH).


- (72) Inventors; and
(75) Inventors/Applicants (*for US only*): HEROLD, Peter [CH/CH]; Unterer Rheinweg 124, CH-4057 Basel (CH). STUTZ, Stefan [CH/CH]; Reichensteinerstrasse 19, CH-4053 Basel (CH).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE PREPARATION OF SUBSTITUTED OCTANOYL AMIDES



(57) Abstract: Compounds of formula (XII), are simultaneously halogenated in the 5 position and hydroxylated in the 4 position under lactonization, the halolactone is reacted with an amine to form a carboxamide, the halogen is replaced with azide, if necessary after the introduction of a hydroxy protecting group, the resulting azide is converted to a lactone, the lactone is amidated and then the azide converted to the amine group, in order to obtain compounds of formula (I) or a salt thereof.

WO 02/08172 A1


Process for the preparation of substituted octanoyl amides

The invention relates to a stereospecific method for the preparation of 2(S),4(S),5(S),7(S)-2,7-dialkyl-4-hydroxy-5-amino-8-aryloctanoyl amides and their physiologically acceptable salts; and new compounds used as intermediates in the multistage process.

In EP-A-0 678 503, δ -amino- γ -hydroxy- ω -aryl-alkanecarboxamides are described, which exhibit renin-inhibiting properties and could be used as antihypertensive agents in pharmaceutical preparations. The manufacturing procedures described are unsatisfactory in terms of the number of process steps and yields and are not suitable for an industrial process. A disadvantage of these processes is also that the total yields of pure diastereomers that are obtainable are too small.

It has now been surprisingly found that these alkanecarboxamides can be prepared both in high total yields and in a high degree of purity, and that selectively pure diastereomers are obtainable, if the double bond of 2,7-dialkyl-8-aryl-4-octenic acid amides is simultaneously halogenated in the 5 position and hydroxylated in the 4 position under lactonization, the lactone ring is opened with an amine during the formation of the carboxamide, then the hydroxy group is replaced with azide, if necessary after protection of the hydroxy group, the resulting compound is lactonized, the lactone amidated and then the azide group converted to the amine group.

A first object of the invention is a process for the preparation of compounds of formula I and their physiologically acceptable salts,




(I),

wherein


R_1 and R_2 are, independently of one another, H, C_1 - C_6 alkyl, C_1 - C_6 halogenalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxy- C_1 - C_6 alkyl, or C_1 - C_6 alkoxy- C_1 - C_6 alkyloxy, R_3 is C_1 - C_6 alkyl, R_4 is C_1 - C_6 alkyl, and R_5 is C_1 - C_6 alkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 alkanoyloxy- C_1 - C_6 alkyl, C_1 - C_6 aminoalkyl, C_1 - C_6 alkylamino- C_1 - C_6 -alkyl, C_1 - C_6 -dialkylamino- C_1 - C_6 -alkyl, C_1 - C_6 -alkanoyl-amido- C_1 - C_6 -alkyl, $HO(O)C$ - C_1 - C_6 -alkyl, C_1 - C_6 alkyl- O -(O) C - C_1 - C_6 alkyl, $H_2N-C(O)$ - C_1 - C_6 alkyl, C_1 - C_6 alkyl- $HN-C(O)$ - C_1 - C_6 alkyl or $(C_1$ - C_6 alkyl) $_2N-C(O)$ - C_1 - C_6 -alkyl, comprising the steps

a) reaction of a compound of formula II,



(II),

with an amine of formula R_5-NH_2 to form a compound of formula III,



(III),


and

b) reduction of the azide group of the compound of formula III to the amine group and isolation of the compounds of

formula I, if necessary with the addition of a salt-forming acid, comprising the preparation of the compound of formula II by reacting

c) a compound of formula IV


5



(IV),

wherein X is Cl, Br or I, with an amine to form a carboxamide of formula V,

10




(V),

wherein R₆ is an amino group,

d1) azidating a compound of formula V to form a compound of


15 formula VI



(VI), or

d2) protecting the hydroxyl group in the compounds of formula V, and azidating the resulting compound of formula VII


20



(VII),

wherein S is a protecting group, to form a compound of formula VIII,

5



(VIII),

e) and then lactonizing the compound of formula VI or VIII in the presence of an acid to form a compound of formula II.

10

As an alkyl, R_1 and R_2 may be linear or branched and preferably comprise 1 to 4 C atoms. Examples are methyl, ethyl, n- and i-propyl, n-, i- and t-butyl, pentyl and hexyl.

15

As a halogenalkyl, R_1 and R_2 may be linear or branched and preferably comprise 1 to 4 C atoms, especially 1 or 2 C atoms. Examples are fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloro-

20

methyl, 2-chloroethyl and 2,2,2-trifluoroethyl.

As an alkoxy, R_1 and R_2 may be linear or branched and preferably comprise 1 to 4 C atoms. Examples are methoxy, ethoxy, n- and i-propyloxy, n-, i- and t-butyloxy, pentyloxy

25

and hexyloxy.

As an alkoxyalkyl, R_1 and R_2 may be linear or branched. The alkoxy group preferably comprises 1 to 4 and especially 1 or

2 C atoms, and the alkyl group preferably comprises 1 to 4 C atoms. Examples are methoxymethyl, 1-methoxyeth-2-yl, 1-methoxyprop-3-yl, 1-methoxybut-4-yl, methoxypentyl, methoxyhexyl, ethoxymethyl, 1-ethoxyeth-2-yl, 1-ethoxyprop-3-yl, 1-ethoxybut-4-yl, ethoxypentyl, ethoxyhexyl, propyloxymethyl, butyloxymethyl, 1-propyloxyeth-2-yl and 1-butyloxyeth-2-yl.

As a C₁-C₆alkoxy-C₁-C₆alkyloxy, R₁ and R₂ may be linear or branched. The alkoxy group preferably comprises 1 to 4 and especially 1 or 2 C atoms, and the alkyloxy group preferably comprises 1 to 4 C atoms. Examples are methoxymethyloxy, 1-methoxyeth-2-yloxy, 1-methoxyprop-3-yloxy, 1-methoxybut-4-yloxy, methoxypentyloxy, methoxyhexyloxy, ethoxymethyloxy, 1-ethoxyeth-2-yloxy, 1-ethoxyprop-3-yloxy, 1-ethoxybut-4-yloxy, ethoxypentyloxy, ethoxyhexyloxy, propyloxymethyloxy, butyloxymethyloxy, 1-propyloxyeth-2-yloxy and 1-butyloxyeth-2-yloxy.

In a preferred embodiment, R₁ is methoxy- or ethoxy-C₁-C₄alkyloxy, and R₂ is preferably methoxy or ethoxy. Particularly preferred are compounds of formula I, wherein R₁ is 1-methoxyprop-3-yloxy and R₂ is methoxy.

As an alkyl, R₃ and R₄ may be linear or branched and preferably comprise 1 to 4 C atoms. Examples are methyl, ethyl, n- and i-propyl, n-, i- and t-butyl, pentyl and hexyl. In a preferred embodiment, R₃ and R₄ in compounds of formula I are in each case isopropyl.

As an alkyl, R₅ may be linear or branched and preferably comprise 1 to 4 C atoms. Examples of alkyl are listed hereinabove. Methyl, ethyl, n- and i-propyl, n-, i- and t-butyl are preferred.

35

As a C₁-C₆hydroxyalkyl, R₅ may be linear or branched and preferably comprise 2 to 6 C atoms. Some examples are 2-hydroxyethyl-1-yl, 2-hydroxyprop-1-yl, 3-hydroxyprop-1-yl, 2-, 3- or 4-hydroxybut-1-yl, hydroxypentyl and hydroxyhexyl.

5

As a C₁-C₆alkoxy-C₁-C₆alkyl, R₅ may be linear or branched. The alkoxy group preferably comprises 1 to 4 C atoms and the alkyl group preferably 2 to 4 C atoms. Some examples are 2-methoxyethyl-1-yl, 2-methoxyprop-1-yl, 3-methoxyprop-1-yl, 2-, 3- or 4-methoxybut-1-yl, 2-ethoxyethyl-1-yl, 2-ethoxyprop-1-yl, 3-ethoxyprop-1-yl, and 2-, 3- or 4-ethoxybut-1-yl.

10

As a C₁-C₆alkanoyloxy-C₁-C₆alkyl, R₅ may be linear or branched. The alkanoyloxy group preferably comprises 1 to 4 C atoms and the alkyl group preferably 2 to 4 C atoms. Some examples are formyloxymethyl, formyloxyethyl, acetyloxyethyl, propionyloxyethyl and butyroyloxyethyl.

15

As a C₁-C₆aminoalkyl, R₅ may be linear or branched and preferably comprise 2 to 4 C atoms. Some examples are 2-aminoethyl, 2- or 3-aminoprop-1-yl and 2-, 3- or 4-aminobut-1-yl.

20

As a C₁-C₆alkylamino-C₁-C₆alkyl and C₁-C₆dialkylamino-C₁-C₆alkyl, R₅ may be linear or branched. The alkylamino group preferably comprises C₁-C₄alkyl groups and the alkyl group preferably 2 to 4 C atoms. Some examples are 2-methylaminoethyl-1-yl, 2-dimethylaminoethyl-1-yl, 2-ethylaminoethyl-1-yl, 2-ethylaminoethyl-1-yl, 3-methylaminoprop-1-yl, 3-dimethylaminoprop-1-yl, 4-methylaminobut-1-yl and 4-dimethylaminobut-1-yl.

25

30

As a C₁-C₆alkanoylamido-C₁-C₆alkyl, R₅ may be linear or branched. The alkanoyl group preferably comprises 1 to 4 C atoms and the alkyl group preferably 1 to 4 C atoms. Some

35

examples are 2-formamidoeth-1-yl, 2-acetamidoeth-1-yl, 3-propionylamidoeth-1-yl and 4-butyroylamidoeth-1-yl.

As a HO(O)C-C₁-C₆alkyl, R₅ may be linear or branched, and the
5 alkyl group preferably comprises 2 to 4 C atoms. Some examples are carboxymethyl, carboxyethyl, carboxypropyl and carboxybutyl.

As a C₁-C₆-alkyl-O-(O)C-C₁-C₆alkyl, R₅ may be linear or
10 branched, and the alkyl groups preferably comprise independently of one another 1 to 4 C atoms. Some examples are methoxycarbonylmethyl, 2-methoxycarbonyleth-1-yl, 3-methoxycarbonylprop-1-yl, 4-methoxycarbonylbut-1-yl, ethoxycarbonylmethyl, 2-ethoxycarbonyleth-1-yl, 3-ethoxycarbonyl-
15 prop-1-yl, and 4-ethoxycarbonylbut-1-yl.

As a H₂N-C(O)-C₁-C₆alkyl, R₅ may be linear or branched, and the alkyl group preferably comprises 2 to 6 C atoms. Some examples are carbamidomethyl, 2-carbamidoeth-1-yl, 2-
20 carbamido-2,2-dimethyleth-1-yl, 2- or 3-carbamidoprop-1-yl, 2-, 3- or 4-carbamidobut-1-yl, 3-carbamido-2-methylprop-1-yl, 3-carbamido-1,2-dimethylprop-1-yl, 3-carbamido-3-methylprop-1-yl, 3-carbamido-2,2-dimethylprop-1-yl, 2-, 3-, 4- or 5-carbamidopent-1-yl, 4-carbamido-3,3- or -2,2-dimethylbut-
25 1-yl.

As a C₁-C₆alkyl-HN-C(O)-C₁-C₆-alkyl or (C₁-C₆alkyl)₂N-C(O)-C₁-C₆-alkyl, R₅ may be linear or branched, and the NH-alkyl group preferably comprises 1 to 4 C atoms and the alkyl
30 group preferably 2 to 6 C atoms. Examples are the carbamidoalkyl groups defined hereinabove, whose N atom is substituted with one or two methyl, ethyl, propyl or butyl.

A preferred subgroup of compounds of formula I is that in
35 which R₁ is C₁-C₄alkoxy or C₁-C₄alkoxy-C₁-C₄alkyloxy, R₂ is C₁-C₄alkoxy, R₃ is C₁-C₄alkyl, R₄ is C₁-C₄alkyl and R₅ is

H₂NC(O)-C₁-C₆alkyl which if necessary is N-monosubstituted or N-di-C₁-C₄alkyl substituted.

5 A more preferred subgroup of compounds of formula I is that in which R₁ is methoxy-C₂-C₄-alkyloxy, R₂ is methoxy or ethoxy, R₃ is C₂-C₄alkyl, R₄ is C₂-C₄alkyl and R₅ is H₂NC(O)-C₁-C₆alkyl.

10 An especially preferred compound of formula I is that in which R₁ is 3-methoxy-prop-3-yloxy, R₂ is methoxy, R₃ and R₄ are 1-methyleth-1-yl, and R₅ is H₂NC(O)-[C(CH₃)₂]-CH₂-.

15 As an amino group, R₆ may be -NH₂, primary and preferably secondary amino, the amino groups comprising 1 to 20 C atoms and preferably 2 to 12. The amino group preferably corresponds to the formula -N(R₇)₂, wherein R₇ is C₁-C₄alkyl, cyclopentyl, cyclohexyl, phenyl or benzyl, or both R₇ are together tetramethylene, pentamethylene or 3-oxapentylene. Preferred examples of R₇ are methyl, ethyl, n-propyl and
20 n-butyl.

Protecting group S in the compounds of formulae VII and VIII are preferably acyl groups, which may comprise 1 to 12 and preferably 1 to 8 C atoms. Some examples are formyl, acetyl,
25 propionyl and butyryl. Acetyl is especially preferred.

The individual process steps may be carried out in the presence of solvent. Suitable solvents are water and organic solvents, especially polar organic solvents, which can also
30 be used as mixtures of at least two solvents. Examples of solvents are hydrocarbons (petroleum ether, pentane, hexane, cyclohexane, methylcyclohexane, benzene, toluene, xylene), halogenated hydrocarbon (dichloromethane, chloroform, tetrachloroethane, chlorobenzene); ether (diethyl ether,
35 dibutyl ether, tetrahydrofuran, dioxane, ethylene glycol dimethyl or diethyl ether); carbonic esters and lactones

(methyl acetate, ethyl acetate, methyl propionate, valerolactone); N,N-substituted carboxamides and lactams (dimethylformamide, dimethylacetamide, N-methylpyrrolidone); ketones (acetone, methylisobutylketone, cyclohexanone);
5 sulfoxides and sulfones (dimethylsulfoxide, dimethylsulfone, tetramethylene sulfone); alcohols (methanol, ethanol, n- or i-propanol, n-, i- or t-butanol, pentanol, hexanol, cyclohexanol, cyclohexanediol, hydroxymethyl or dihydroxymethyl cyclohexane, benzyl alcohol, ethylene glycol,
10 diethylene glycol, propanediol, butanediol, ethylene glycol monomethyl or monoethyl ether, and diethylene glycol monomethyl or monoethyl ether; nitriles (acetonitrile, propionitrile); tertiary amines (trimethylamine, triethylamine, tripropylamine and tributylamine, pyridine, N-methylpyrrolidine,
15 pyrrolidine, N-methylpiperazine, N-methylmorpholine) and organic acids (acetic acid, formic acid).

Process step a)

The reaction of compounds of formula II with a compound R_5NH_2 by opening of the lactone ring to form compounds of formula
20 III is expediently carried out in the presence of alcohols or amines which are capable of forming activated carbonic esters or carboxamides. Such compounds are well-known. They may be 2-hydroxypyridine, N-hydroxycarboxamides and imides,
25 and carboximides (N-hydroxysuccinimide). Organic solvents are used as solvent, tertiary amines being of advantage, for example trimethylamine or triethylamine. The reaction temperature may range for example from approximately 40°C to 150°C and preferably from 50°C to 120°C.

30

Process step b)

Reduction of the azide group to the amine group in the compounds of formula III takes place in a manner known per se (see Chemical Reviews, Vol. 88 (1988), pages 298 to 317),
35 for example using metal hydrides or more expediently using a catalytic method with hydrogen in the presence of

homogeneous (Wilkinson catalyst) or heterogeneous catalysts, for example Raney nickel or precious metal catalysts such as platinum or palladium, if necessary on substrates such as carbon. The hydrogenation can also be carried out if
5 necessary catalytically under phase transfer conditions, for example with ammonium formate as hydrogen donor. It is of advantage to use organic solvents. The reaction temperature may range for example from approximately 0°C to 200°C and preferably from 10°C to 100°C. Hydrogenation may be carried
10 out at normal pressure or increased pressure up to 100 bar, for example, and preferably up to 50 bar.

The compounds of formula I may be converted to addition salts in a manner known per se by treatment with monobasic
15 or polybasic, inorganic or organic acids. Hemifumarates are preferred.

Process step c)

The reaction of the halolactone with an amine to form
20 carboxamide is advantageously carried out in organic solvents such as halogenated hydrocarbons (chloroform, dichloromethane). The reaction temperature may range for example from approximately -30°C to 80°C and preferably from -20°C to 50°C. The amine is expediently used as a salt, for
25 example as a halogenide. Dimethyl ammonium chloride is preferably used. The reaction is preferably carried out in the presence of at least equimolar quantities of an alkyl aluminium halogenide such as dialkyl aluminium chloride (dimethyl or diethyl aluminium chloride). After hydrolytic
30 treatment, the carboxamide can be isolated by means of extraction and purified by means of chromatography. The stereoselectivity is high and the yield can be as much as 70% or more.

Process step d1)

Halogen X may be directly substituted with azide in the carboxamide of formula V obtained as described in step c). Suitable azidation agents are for example metal azides, especially alkaline earth metal azides and alkali metal azides, as well as silyl azides. Especially preferred azidation agents are lithium azide, sodium azide and potassium azide. The reaction may be carried out in organic solvents, for example N-alkylated lactams such as N-methylpyrrolidone or 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone (DMPU). The reaction temperature may range for example from approximately 20°C to 150°C and preferably from 20°C to 120°C. It may be expedient to include the use of phase transfer catalysts. In the broader sense it is advantageous to carry out the reaction in the presence of preferably at least equimolar quantities of a base, especially tertiary amines. These tertiary amines may serve at the same time as solvents. The preparation and synthetic use of azides are described for example by E. F. V. Scriven in Chemical Reviews, Vol. 88 (1988), pages 298 to 317. As a result of secondary reactions due to the absence of the hydroxyl group, the yield in the non-optimized reaction is not very high and may be about 30% or more.

25 Process step d2)

It has therefore proved very advantageous to protect the hydroxyl group against azidation in the compounds of formula VI, preferably with acyl groups. To this end, compounds of formula V are reacted with acylation agents, for example carboxylic acid anhydrides such as acetic acid anhydride or carboxylic acid halogenides such as acetylchloride. The reaction may be carried out with or without solvents. The reaction temperature may be -20 to 80°C. The reaction is expediently carried out in the presence of bases, for example tertiary amines. Examples of tertiary amines are trialkylamines (trimethylamine, triethylamine), N-alkylated

cyclic amines (N-alkylpyrrolidine), dialkylaminopyridines (dimethylaminopyridine) and pyridine. After hydrolytic treatment, the protected carboxamide can be isolated by means of extraction and purified by means of chromatography.

5 The yield is generally more than 90%.

Azidation may then be carried out as described in process step d1). The yields are substantially higher than with direct azidation as described in process step d1) and are

10 more than 75% in the non-optimized process step d2).

Process step e)

Lactonization of compounds of formula VI or VIII to form compounds of formula II is expediently carried out at a


15 temperature of -20 to 100°C and in the presence of a solvent such as alcohols (methanol, ethanol or propanol) or hydrocarbons (benzene, toluene or xylene). Inorganic acids and advantageously organic acids are used, especially mineral acids such as hydrochloric acid, hydrobromic acid or

20 sulfuric acid, sulfonic acids and carboxylic acids. The azidolactone of formula II may be isolated for example by extraction with organic solvents. The desired stereoisomer is also formed in this step at high yields of up to 90% or more.

25

Some intermediates prepared using the process according to the invention are new and represent further objects of the invention.


30 A further object of the invention is thus a compound of formula IX,



(IX),

wherein

- X is halogen, R_1 and R_2 are, independently of one another, H, C_1 - C_6 alkyl, C_1 - C_6 -halogenalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxy- C_1 - C_6 alkyl, or C_1 - C_6 alkoxy- C_1 - C_6 alkyloxy, R_3 is C_1 - C_6 alkyl, R_4 is C_1 - C_6 alkyl, R_6 is an amino group, and R_8 is a protecting group or hydrogen.
- 10 A further object of the invention is a compound of formula IXa,




(IXa),

- 15 wherein

R_1 and R_2 are, independently of one another, H, C_1 - C_6 alkyl, C_1 - C_6 -halogenalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxy- C_1 - C_6 alkyl, or C_1 - C_6 alkoxy- C_1 - C_6 alkyloxy, R_3 is C_1 - C_6 alkyl, R_4 is C_1 - C_6 alkyl, R_6 is an amino group, and R_8 is a protecting group or hydrogen.


20 For residues X, R_1 , R_2 , R_3 , R_4 , R_6 , and R_8 in compounds of formulae IX and IXa, the embodiments and preferences described hereinbefore apply.

- 25 The compounds of formula IV are obtainable by reacting in a first step a compound of formula X,



(X),

with a compound of formula IX,




5

(XI),

wherein R_1 to R_4 are as defined hereinbefore, including the preferences, Y is Cl , Br or I and Z is Cl , Br or I (Y and Z are preferably Br and especially Cl), and R_9 is C_1 - C_6 alkyl, R_{10} is C_1 - C_6 -akyl or C_1 - C_6 alkoxy, or R_9 and R_{10} are together tetramethylene, pentamethylene, 3-oxa-1,5-pentylene or - $CH_2CH_2O-C(O)-$ substituted if necessary with C_1 - C_4 alkyl, phenyl or benzyl, in the presence of an alkali or alkaline earth metal to form a compound of formula XII,

15



(XII),

wherein

R_9 is C_1 - C_6 alkyl, R_{10} is C_1 - C_6 alkyl or C_1 - C_6 alkoxy, or R_9 and R_{10} together are tetramethylene, pentamethylene, 3-oxa-1,5-pentylene or - $CH_2CH_2O-C(O)-$ substituted if necessary with C_1 - C_4 alkyl, phenyl or benzyl.

As an alkyl, R_9 and R_{10} in formula XII may be branched and preferably linear and are preferably C_1 - C_4 alkyl, for example methyl or ethyl. R_{10} as alkoxy may preferably be linear and is preferably C_1 - C_4 alkoxy, for example methoxy or ethoxy.

R₉ and R₁₀ together are preferably tetramethylene, -CH₂CH₂-O-C(O)- or -CH(CH₂C₆H₅)CH₂-O-C(O)-.

The coupling of Grignard reagents with alkenyl halogenides
5 in an ether such as, for example, tetrahydrofuran or dioxan
as solvents in the presence of catalytic quantities of a
soluble metal complex, for example an iron complex such as
iron acetylacetonate, and in the presence of more than
equimolar quantities of a solvent stabilizing the metal
10 complex, for example N-methylpyrrolidone, is described by G.
Cahiez et al. in Synthesis (1998), pages 1199-1200. The
reaction temperature may for example be -50 to 80°C,
preferably -20 to 50°C. Catalytic quantities may for example
be 0.1 to 20% by weight in relation to a compound of formula
15 VII. It is expedient to carry out the reaction so that
initially a compound of formula VI is converted to a
Grignard compound (for example with magnesium) and then
adding a solution of a compound of formula VII, metal
complex and N-methylpyrrolidone, or vice versa.

20

It was found to be of advantage when only catalytic
quantities of a solvent stabilizing the metal complexes, for
example N-methylpyrrolidone, were used. Catalytic quantities
may for example be 1 to 10 mol per cent, preferably 1 to 5
25 mol per cent, in relation to the compounds of formula XI or
XII.

Compounds of formula X in the form of their racemates or
enantiomers are known or capable of being prepared according
30 to analogous processes. For example, R₁R₂phenylaldehyde may
be reacted with R₃diethoxyphosphorylacetic acid ester to form
2-R₃-3-(R₁R₂phenyl)acrylic acid esters, these may then be
hydrogenated to form the corresponding propionic acid
esters, the ester group saponified and the carboxylic acid
35 reduced to alcohol, and finally the hydroxyl group
substituted with halogen. Enantiomers are obtainable by

separating the racemates of the carboxylic acids with for example quinine or by enzymatic resolution of the racemates of the corresponding carboxylic acid esters. Details are described in the examples. A possible asymmetric synthesis
5 of compounds of formula VI is described in EP-A-0 678 503.

Compounds of formula XI in the form of their racemates or enantiomers may be prepared by the reaction of metallated carbonic esters of formula R_4CH_2COOR (for example lithium
10 isovaleric acid esters) with trans-1,3-halogenpropene, then halogenation of the resulting carboxylic acid to form the acid halogenide and reaction with a secondary amine. The coupling of metallated carbonic esters with trans-1,3-halogenpropene can be carried out asymmetrically according
15 to the method described by D. A. Evans in Asymmetric Synthesis, Vol. 3, 1984 (Academic Press Inc.), pages 2-110. Enantiomers are obtainable by separating the racemates of the carboxylic acids with for example cinchonidine or by enzymatic separation of the racemates of the corresponding
20 carbonic esters.

In a second process step, compounds of formula XII are reacted with a halogenation agent in the presence of water and if necessary an acid to form a compound of formula IV.
25

Suitable chlorination, bromination and iodination agents are elemental bromine and iodine, in particular N-chloro, N-bromo and N-iodocarboxamides and dicarboximides. Preferred are N-chloro, N-bromo and N-iodophthalimide and especially
30 chloro, N-bromo and N-iodosuccinimide, as well as tertiary butyl hypochlorite and N-halogenated sulfonamides and sulfonimides, for example chloramine T. It is of advantage to carry out the reaction in organic solvents. The reaction temperature may range for example from approximately $-70^{\circ}C$
35 to ambient temperature and preferably from $-30^{\circ}C$ to $10^{\circ}C$. Carboxamides are advantageously lactonized in the presence

of inorganic or organic acids, at least equimolar quantities of water, and reacted in the presence of water-miscible solvents, for example tetrahydrofuran or dioxane. Suitable acids are for example formic acid, acetic acid, 5 methanesulfonic acid, trifluoroacetic acid, trifluoromethanesulfonic acid, toluenesulfonic acid, H_2SO_4 , H_3PO_4 , hydrogen halides, acid ion exchange resins, and acids immobilized on solid carriers. Water is generally used in at least equimolar quantities.


10

With the choice of lactones of formula IV, the compounds of formula I, which per se are complex compounds, can be prepared in a convergent and simple manner, which is especially true of these enantioselective or diastereo- 15 selective synthesis. The total yield from all process steps a) to e) may amount to 40% or more, which makes an industrial application feasible.


The following examples explain the invention in more detail.

20

A) Preparation of compounds of formula X



(EP-A-0 678 503)




(A1)

25 Example A1:

An agitated solution of 174 g 2R-[4-methoxy-3-(3-methoxypropoxy)benzyl]-3-methylbutan-1-ol [EP 0 678 503] and 1.3 l carbon tetrachloride is cooled to $10^\circ C$. 393 ml trioctylphosphine is added dropwise, and the reaction 30 solution is then stirred for 16 hours at ambient temperature. The mixture is completely concentrated by evaporation and the residue extracted between dichloromethane (3x) and water (1x). The combined organic

phases are dried over magnesium sulfate, filtered and concentrated by evaporation. The residue is purified by means of flash chromatography (SiO₂ 60F / ethyl acetate / hexane 1:9), and title compound A5 is obtained after
5 crystallization (hexane at -°50C) as a white solid (152.3 g, 82 %): melting point 51-52°C; ¹H-NMR (400 MHz, CDCl₃, δ): 1.0 (m, 6H), 1.71 (m, 1H), 1.93 (m, 1H), 2.12 (m, 2H), 2.35 (m, 1H), 2.77 (m, 1H), 3.39 (s, 3H), 3.40 - 3.55 (m, 2H), 3.71 (t, 2H), 3.87 (s, 3H), 4.13 (m, 3H), 6.65 - 6.85 (m, 3H)
10 ppm.

B) Preparation of compounds of formula XI



Example B1:

An agitated solution of 24.9 ml diisopropylamine and 240 ml tetrahydrofuran is cooled to minus 15°C, and 100 ml 1.6 M n-butyl lithium solution (in hexane) is added over a period of 10 minutes. The solution is stirred for 30 minutes at -15°C and then, over a period 30 minutes, a solution of 24.1 ml ethyl isovalerate in 80 ml tetrahydrofuran is added dropwise. The mixture is stirred for a further 5 minutes at -15°C, and then 19.5 g trans-1,3-dichloropropene and 2.4 g

sodium iodide are added consecutively. The reaction mixture is stirred for a further 16 hours at ambient temperature, and then 500 ml 10% aqueous ammonium chloride solution is added. The mixture is extracted with diethyl ether (3x) and the organic phases washed consecutively with water (1x), 0.1 M sodium thiosulfate solution (1x) and brine (1x). The combined organic phases are dried with sodium sulfate and concentrated by evaporation. By means of distillation, title compound B1 is obtained as a colourless oil (24.8 g, 76 %).

10 $^1\text{H-NMR}$ (400 MHz, CDCl_3 , δ): 0.95 (m, 6H), 1.30 (t, 3H), 1.92 (m, 1H), 2.20 - 2.40 (m, 3H), 4.20 (m, 2H), 5.80 - 6.10 (m, 2H) ppm.

Example B2:

15 A solution of 150.2 g B1, 500 ml ethanol and 500 ml 2N sodium hydroxide solution is stirred for 18 hours under reflux. The ethanol is evaporated from the reaction mixture, the aqueous solution acidified with 1N hydrochloric acid and extracted with diethyl ether (3x). The organic phases are dried over magnesium sulfate and concentrated by evaporation. By means of flash chromatography (SiO_2 60F / dichloromethane / methanol 20:1), title compound B2 is obtained from the residue as a slightly orange oil (83.7 g, 65 %): $^1\text{H-NMR}$ (400 MHz, CDCl_3 , δ): 1.03 (m, 6H), 1.98 (m, 1H), 2.20 - 2.45 (m, 3H), 5.80 - 6.10 (m, 2H) ppm.

25

Example B3: Racemate resolution of compound B2

5.0 g B2, 5.0 g cinchonidine and 1.98 ml triethylamine are transferred to 150 ml tetrahydrofuran and stirred for 15 minutes under reflux. The oil bath is removed and the clear solution with a salt of B3 is inoculated with cinchonidine. Agitation is continued for 1 hour at ambient temperature and then for another 1 hour under ice cooling. The precipitate is filtered off, washed with twice 25 ml ice-cold acetone and then dried in a vacuum at 50°C until constant weight is attained. 6.16 g (46.3%) of the enriched salt of B3 is

35

obtained with cinchonidine; melting point 149°C. After double recrystallization from acetone, 4.20 g (31.6%) of the enriched salt of B3 is obtained with cinchonidine, melting point 155°C. The salt obtained in this way is distributed
5 between 250 ml diethyl ether and 50 ml 1N HCl. The aqueous phase is separated off, the organic phase washed with saturated NaCl solution, dried over MgSO₄ and concentrated by evaporation in a vacuum. 1.58 g (31.6%) of enriched compound B3 is obtained as colourless oil.

10

Example B4: Asymmetric synthesis of B3

a) A solution of 290 g 4S-benzyl-3-(3-methyl-butyl)oxazolidin-2-one in 0.58 l tetrahydrofuran is cooled to -78°C, and 1.14 l 1 M lithium hexamethyldisilazide (in
15 tetrahydrofuran) is added dropwise over a period of 65 minutes. The mixture is stirred for another hour at -78°C, and a prepared solution of trans-1-chloro-3-iodopropene in tetrahydrofuran is then added. The temperature is allowed to increase to 0°C and agitation is continued for a further 20
20 hours. 500 ml 10% ammonium chloride solution is added to the reaction mixture, which is then extracted with diethyl ether (2x 1 l). The organic phases are washed with water (1x 1 l), dried with sodium sulfate and concentrated by evaporation. By means of flash chromatography (SiO₂ 60F / ethyl acetate /
25 hexane 5:1), title compound B5 is obtained from the residue as a slightly orange oil (582 g, 78 %): ¹H-NMR (400 MHz, CDCl₃, δ): 0.85 (m, 6H), 2.02 (m, 1H), 2.3 - 2.55 (m, 2H), 2.75 (m, 1H), 3.30 (m, 1H), 3.88 (m, 1H), 4.18 (m, 2H), 4.70 (m, 1H), 5.80 - 6.10 (m, 2H), 7.15 - 7.40 (m, 5H) ppm.

30 Racemate resolution of compound A3 266.1 g sodium iodide is added to a solution of 184.7 g trans-1,3-dichloropropene in 0.58 l tetrahydrofuran and the mixture stirred for 30 minutes under exclusion of light at ambient temperature. The mixture is filtered until clear and the filtrate used in the
35 crude state.


b) To a solution of 155 g B4, 1.3 l tetrahydrofuran and 0.44 l water, stirred at 0°C, 315 ml 30% hydrogen peroxide solution is added dropwise over a period of 15 minutes. 22.1 g lithium hydroxide is added to the reaction mixture, then the cooling bath is removed and stirring is continued for 5 hours at 0-20°C. The reaction mixture is cooled again to 0°C, and a solution of 350 g sodium sulfite in 1.4 l water is added dropwise over a period of 30 minutes. The pH is adjusted to 9.8 by the addition of sodium hydrogencarbonate. The reaction mixture is filtered until clear and tetrahydrofuran evaporated from the filtrate. The aqueous solution obtained is washed with dichloromethane (3x 3 l). The pH of the aqueous phase is adjusted to 3.0 with aqueous hydrochloric acid and then extracted with dichloromethane (3x 2l). The organic phases are dried over magnesium sulfate and concentrated by evaporation on a rotary evaporator. By means of distillation, title compound B3 is obtained from the residue as a colourless oil. (142 g, 87 %). ¹H-NMR (400 MHz, CDCl₃, δ): 1.02 (m, 6H), 1.98 (m, 1H), 2.25 - 2.45 (m, 3H), 5.85 - 6.10 (m, 2H) ppm.

Example B4:

4.42 ml oxalyl chloride is added to a solution of 4.54 g B3 in 25 ml toluene at ambient temperature. The reaction mixture is agitated for 15 minutes at ambient temperature, and then 0.052 ml N,N-dimethylformamide over a period of 1 minute. The reaction mixture is heated to reflux and agitated for 1 hour. The reaction solution is concentrated by evaporation and the residue distilled. Title compound B4 is obtained as a colourless oil. (4.43 g, 88 %). ¹H-NMR (400 MHz, CDCl₃, δ): 1.02 (d, 3H), 1.08 (d, 3H), 2.16 (m, 1H), 2.40 (m, 1H), 2.45 (m, 1H), 2.68 (m, 1H), 5.80 - 6.10 (m, 2H) ppm.

Example B5:

A solution of 1.53 g dimethylamine, 3.66 ml pyridine and 25 ml dichloromethane is cooled to 0°C, and then 4.42 g B5 in 25 ml dichloromethane is added dropwise at 0 to -10°C. The reaction mixture is stirred for a further 2 hours at 0°C and then concentrated by evaporation on the Rotavapor. The residue is distributed between diethyl ether (2x) and 2N hydrochloric acid (3x), saturated sodium hydrogencarbonate solution (1x) and saturated saline solution. The organic fractions are combined, dried over sodium sulfate and concentrated. The residue is distilled, and title compound B6 is obtained as a colourless oil. (4.13 g, 89 %). $[\alpha]_D^{25}$ -7,3 (c 1, chloroform). $^1\text{H-NMR}$ (400 MHz, CDCl_3 , δ): 0.90 (d, 3H), 0.95 (d, 3H), 1.92 (m, 1H), 2.20 - 2.30 (m, 1H), 2.35 - 2.50 (m, 2H), 2.98 (s, 3H), 3.04 (s, 3H), 5.80 - 6.10 (m, 2H) ppm.


Example B6:

A mixture of 10.7 g magnesium powder and 120 ml tetrahydrofuran is heated to 60°C, and 0.74 ml 1,2-dibromoethane then added over a period of 2 minutes (visible exothermic reaction). A solution of 34.6 g A1, 4.0 ml 1,2-dibromoethane and 320 ml tetrahydrofuran is added dropwise over a period of 15 minutes at 62 - 64°C. The mixture is stirred for another 30 minutes under reflux and then cooled down to ambient temperature. The reaction mixture is filtered under argon until clear and the resulting Grignard solution added dropwise over a period of 10 minutes to a solution of 20.4 g B6, 0.240 ml N-methylpyrrolidone, 0.88 g iron(III) acetylacetonate in 200 ml tetrahydrofuran at -5 to 0°C. The reaction mixture is agitated for a further 15 minutes at 0 to 10°C, and 320 ml

2N hydrochloric acid is then added. The mixture is now extracted with diethyl ether (3x 500 ml) and the organic phases washed consecutively with water (1x 400 ml) and saturated aqueous sodium chloride solution (1x 400 ml). The combined organic phases are dried over sodium sulfate, filtered and concentrated on a rotary evaporator. By means of flash chromatography (SiO₂ 60F / diethyl ether / hexane 2:1), title compound B7 is obtained from the residue as a slightly yellowish oil (36.2 g, 81 %): TLC R_t = 0.09 (diethyl ether / hexane 2:1); ¹H-NMR (500 MHz, CDCl₃, δ): 0.82 - 0.99 (m, 12H), 1.49 (m, 1H), 1.69 (m, 1H), 1.78 - 1.98 (m, 3H), 2.10 (m, 2H), 2.17 - 2.41 (m, 5H), 2.92 (s, 3H), 3.0 (s, 3H), 3.37 (s, 3H), 3.58 (t, 2H), 3.84 (s, 3H), 4.10 (t, 2H), 5.26 - 5.34 (m, 1H), 5.36 - 5.44 (m, 1H), 6.64 (m, 2H), 6.78 (d, 1H) ppm.

C) Preparation of compounds of formula IV

Example C1: Preparation of




(C1)

3.85 ml water is added to a solution of 34.2 g B7 and 385 ml tetrahydrofuran, and the mixture cooled to 0°C while being stirred. Then 10 times 1.03 ml 42.5 % o-phosphoric acid and 10 times 1.5 g N-bromosuccinimide are added alternately every 3 minutes. The reaction mixture is agitated for another 90 minutes at 0°C and then, over a period of 10 minutes, is introduced to 600 ml sodium hydrogen sulfite solution cooled to 0°C. The mixture is agitated for another 15 minutes at 0°C and then extracted with diethyl ether (1x 1 l and 2x 0.5 l). The organic phases are washed

consecutively with 1N hydrochloric acid (1x 0.6 l), water (1x 0.6 l), saturated aqueous sodium hydrogencarbonate solution (1x 0.6 l) and brine (1x 0.6 l), dried over sodium sulfate and concentrated by evaporation on a rotary
5 evaporator. By crystallization (diisopropyl ether-hexane 1:2 at -25°C), title compound C1 is obtained as a white crystallizate (27.5 g, 72 %): Melting point 48 -49°C; TLC R_f = 0.09 (diethyl ether / hexane 2:1); $[\alpha]_D^{25} = 44.2$ (c 1, chloroform); $^1\text{H-NMR}$ (500 MHz, CDCl_3 , δ): 0.85 - 1.07 (m, 12H), 1.57 -1.65 (m, 1H), 1.79 -2.00 (m, 3H), 2.07 -2.27 (m, 6H), 2.62 (m, 1H), 2.75 (dd, 1H), 3.37 (s, 3H), 3.59 (t, 2H), 3.86 (s, 3H), 4.02 (m, 1H), 4.12 (t, 2H), 4.35 (m, 1H), 6.72 (dd, 1H), 6.75 (d, 1H), 6.81 (d, 1H) ppm.

15 D) Preparation of compounds of formula V




Example D1:

(D1)

A mixture of 6.52 g dimethylamine hydrochloride and 400 ml dichloromethane is cooled to -4°C, and 44.8 ml diethyl-
20 aluminium chloride (1.8M in toluene) is added over a period of 10 minutes. The temperature is allowed to rise to 20°C, a solution of 20 g C1 in 80 ml dichloromethane is added and the mixture stirred for another 18 hours at 35°C. The reaction solution is cooled to 0°C and then stirred in drop
25 by drop to 800 ml 0.5N cold hydrochloric acid. The reaction mixture is extracted with tertiary butyl methyl ether (2x 250 ml), and the resulting organic phases are consecutively washed with water (500 ml) and (concentrated aqueous saline solution (brine, 200 ml). The combined organic phases are
30 dried over sodium sulfate, filtered and concentrated by evaporation. By means of flash chromatography (SiO_2 60F / ethyl acetate / hexane 1:1), title compound D1 is obtained

from the residue as a slightly yellowish oil (19.0 g, 68 %):
 TLC R_t = 0.16 (ethyl acetate / hexane 1:1); $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 0.70 - 0.95 (m, 12H), 1.50 - 1.95 (m, 7H), 2.05 (m, 2H), 2.20 (m, 1H), 2.55 - 2.80 (m, 3H), 2.90 (s, 3H),
 5 3.05 (s, 3H), 3.30 (s, 3H), 3.45 (m, 1H), 3.50 (t, 2H), 3.80 (s, 3H), 4.05 (t, 2H), 4.15 (m, 1H), 6.60 - 6.75 (m, 3H) ppm.


E) Preparation of compounds of formula VII



10 Example E1:

(E1)


A solution of 8.30 g D1 in 100 ml dichloromethane is mixed with 1.54 ml pyridine and cooled to 0°C. Then 1.73 ml acetic acid anhydride and 0.186 g 4-dimethylaminopyridine are added consecutively and the mixture is stirred for 18 hours at
 15 room temperature. The reaction mixture is poured onto 300 ml water and extracted with diethyl ether (2x 300 ml). The organic phases are washed consecutively with water (300 ml), 5% aqueous sodium hydrogencarbonate solution (100 ml) and brine (100 ml). The combined organic phases are dried over
 20 sodium sulfate and concentrated by evaporation on a rotary evaporator. By means of flash chromatography (SiO_2 60F / diethyl ether / hexane 1:1), title compound E1 is obtained from the residue as a colourless oil (7.67 g, 92 %): TLC R_t = 0.27 (ethyl acetate / hexane 1:1); $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ
 25 0.80 - 1.00 (m, 12H), 1.65 - 2.20 (m, 9H), 2.10 (s, 3H), 2.35 (m, 1H), 2.50 - 2.65 (m, 2H), 3.00 (d, 6H), 3.40 (s, 3H), 3.60 (t, 2H), 3.85 (s, 3H), 4.15 (t, 2H), 4.10 (m, 1H), 4.70 (m, 1H), 6.70 - 6.85 (m, 3H) ppm.

F) Preparation of compounds of formula VIExample F1:

(F1)

- 5 A mixture of 0.10 g D1, 0.024 g sodium azide and 1 ml DMPU is stirred for 96 hours at 30°C. The reaction mixture is cooled to room temperature, 30 ml water added, and extraction carried out using diethyl ether (2x 30 ml). The combined organic phases are washed with water (2 x 30 ml)
- 10 and brine (1 x 10 ml), dried over sodium sulfate, filtered and concentrated by evaporation. By means of flash chromatography (SiO₂ 60F / ethyl acetate / hexane 1:1), title compound F1 is obtained from the residue as a colourless oil (27 mg, 29 %): TLC R_t = 0.14 (ethyl acetate / hexane 1:1);
- 15 ¹H-NMR (300 MHz, CDCl₃): δ 0.75 - 0.90 (m, 12H), 1.10 -1.95 (m, 7H), 2.05 (m, 2H), 2.45 (d, 2H), 2.55 (d, 1H), 2.70 (m, 1H), 2.80 - 2.95 (m, 1H), 2.95 (s, 3H), 3.05 (s, 3H), 3.20 - 3.35 (m, 1H), 3.30 (s, 3H), 3.50 (t, 2H), 3.80 (s, 3H), 4.05 (t, 2H), 6.60 -6.75 (m, 3H) ppm.

20


G) Preparation of compounds of formula VIIIExample G1:

(G1)

- 25 A mixture of 1.17 g E1, 0.392 g lithium azide and 11.7 ml DMPU is stirred for 21 h at 60°C. The reaction mixture is cooled, and water (100 ml) added. Extraction is carried out using tertiary butyl methyl ether (3 x 80 ml) and the organic phases are then washed consecutively with water

(3 x 100 ml), 5% aqueous sodium hydrogencarbonate solution (100 ml) and brine (100 ml). The combined organic phases are dried over sodium sulfate and concentrated by evaporation on a rotary evaporator. By means of flash chromatography (SiO₂ 60F / diethyl ether / hexane 3:1), title compound G1 is obtained from the residue as a colourless oil (0.83 g, 76 %): TLC R_t = 0.06 (diethyl ether / hexane 3:1); ¹H-NMR (300 MHz, CDCl₃): δ 0.80 - 1.00 (m, 12H), 1.15 - 1.20 (m, 1H), 1.40 - 2.20 (m, 8H), 2.05 (s, 3H), 2.40 - 2.60 (m, 3H), 3.00 (d, 6H), 3.05 (m, 1H), 3.40 (s, 3H), 3.60 (t, 2H), 3.90 (s, 3H), 4.15 (t, 2H), 4.75 (m, 1H), 6.70 - 6.85 (m, 3H) ppm.


H) Preparation of compounds of formula II



Example H1:


(H1)

A mixture of 70 mg F1, 2 ml toluene and 0.16 ml ethyl acetate is stirred for 4 hours at room temperature. The reaction mixture is cooled to room temperature, 5% aqueous sodium hydrogencarbonate solution (25 ml) is added, and extraction carried out using diethyl ether (2x 20 ml). The combined organic phases are dried over sodium sulfate and concentrated by evaporation on a rotary evaporator. The dried residue corresponds to crude title compound H1 (quantitative); TLC R_t = 0.41 (ethyl acetate / hexane 1:1); ¹H-NMR (300 MHz, CDCl₃): δ 0.85 - 1.10 (m, 12H), 1.40 (m, 1H), 1.60 - 2.25 (m, 8H), 2.45 (m, 1H), 2.60 (m, 2H), 2.95 (m, 1H), 3.40 (s, 3H), 3.60 (t, 2H), 3.85 (s, 3H), 4.15 (t, 2H), 4.30 (m, 1H), 6.70 - 6.85 (m, 3H) ppm.

Example H2:

(H1)

A mixture of 55 mg G1, 38 mg p-toluenesulfonic acid hydrate and 1 ml methyl alcohol is stirred under reflux for 16 hours. The reaction mixture is cooled to room temperature, 5% aqueous sodium hydrogencarbonate solution (5 ml) is added, and extraction carried out using diethyl ether (2x 10 ml). The combined organic phases are dried over sodium sulfate and concentrated by evaporation on a rotary evaporator. The dried residue corresponds to crude title compound H1 (0.043 g, 93%); TLC R_f = 0.41 (ethyl acetate / hexane 1:1); $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 0.85 - 1.10 (m, 12H), 1.40 (m, 1H), 1.60 - 2.25 (m, 8H), 2.45 (m, 1H), 2.60 (m, 2H), 2.95 (m, 1H), 3.40 (s, 3H), 3.60 (t, 2H), 3.85 (s, 3H), 4.15 (t, 2H), 4.30 (m, 1H), 6.70 - 6.85 (m, 3H) ppm.


J) Preparation of compounds of formula IIIExample J1: Preparation of

A mixture of 59.1 g H1, 41.82 g 3-amino-2,2-dimethylpropionamide, 2.28 g 2-hydroxypyridine in 59.1 ml triethylamine is stirred for 16 hours at 90°C. Then 33 ml triethylamine is distilled off over a period of 0.5 hours, and the residue is agitated for a further 8.5 hours at 90°C. The cooled reaction mixture is extracted between ethyl acetate (3x 500 ml), saturated aqueous sodium hydrogencarbonate solution

(1x 500 ml) and saturated sodium chloride solution (1x 500 ml). The combined organic phases are dried over 100 g sodium sulfate, filtered and concentrated on a rotary evaporator. The residue is dried and crude title compound F1 is obtained
5 as an oil (78.4g, quantitative) (HPLC assay: 88.5%): TLC R_t = 0.13 (ethyl acetate - hexane 4:1); chromatographed sample: TLC R_t = 0.13 (ethyl acetate / hexane 4:1); $^1\text{H-NMR}$ (500 MHz, CDCl_3 , δ): 0.85 - 0.96 (m, 12H), 1.23 (s, 6H), 1.30 - 1.40 (m, 1H), 1.53 - 1.80 (m, 5H), 1.82 - 1.93 (m, 1H), 2.06 -
10 2.14 (m, 3H), 2.45 - 2.57 (m, 2H), 2.87 - 2.92 (m, 1H), 3.13 (d, 1H), 3.32 - 3.52 (m, 3H), 3.36 (s, 3H), 3.59 (t, 2H), 3.84 (s, 3H), 4.12 (t, 2H), 5.51 (bs, 1H), 6.01 (bs, 1H), 6.43 (t, 1H), 6.72 (dd, 1H), 6.75 (d, 1H), 6.81 (d, 1H) ppm.

15 K) Hydrogenation of the azide group to form compounds of formula I

Example K1: Preparation of




78.4 g (HPLC assay: 88.5%) F1 (crude) is hydrogenated for 3 hours in the presence of 3.92 g Pd/C 5% and 7.2 ml ethanol amine in 700 ml tert-butyl methyl ether at ambient
25 temperature and 3.0 bar. The reaction mixture is filtered and the catalyst washed with 300 ml tert-butyl methyl ether. The filtrate is washed consecutively with 400 ml 2N NaOH and 400 ml brine. The aqueous phases are then extracted with tert-butyl methyl ether (2 x 400 ml). The combined organic
30 phases are dried over 100 g sodium sulfate and concentrated by evaporation. The residue is mixed with 7.31 g fumaric acid and dissolved in 200 ml ethanol and filtered until

clear. The filtrate is concentrated by evaporation to a total weight of 104 g and dissolved in 1.7 l acetonitrile at 35°C. The resulting solution is inoculated with 10 mg of title compound (hemifumarate) and agitated for 17 hours at ambient temperature. The suspension is cooled to 0°C and filtered off by suction after 2 hours. The residue is washed with acetonitrile (3 x 200 ml) and then dried in a vacuum at 35°C. The title compound K1 (hemifumarate) is obtained as white crystals (59.5 g, 81% in relation to J1):-{}- ¹H NMR (360 MHz, DMSO-d₆); δ 0.7 - 0.9 (m, 12H), 1.04 (s, 6H), 1.27 (m, 3H), 1.4 - 1.8 (m, 4H), 1.94 (m, 2H), 2.23 (m, 1H), 2.35 (dd, J = 8.4, 8.0 Hz, 1H), 2.45 (m, 1H), 3.08 (m, 2H), 3.2 - 3.5 (m, 2H), 3.24 (s, 3H), 3.47 (t, J = 6.4 Hz, 2H), 3.74 (s, 3H), 3.97 (t, J = 6.4 Hz, 2H), 6.37 (s, 1H), 6.68 (dd, J = 8.0, 2.0 Hz, 1H), 6.77 (d, J = 6 Hz, 1H), 6.80 (bs, 1H), 6.83 (d, J = 8 Hz, 1H), 7.13 (bs, 1H), 7.49 (t, J = 6 Hz, 1H).

What is claimed is:

1. A process for the preparation of compounds of formula I and their physiologically acceptable salts,


5



(I),

wherein


10 R₁ and R₂ are, independently of one another, H, C₁-C₆alkyl, C₁-C₆halogenalkyl, C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, or C₁-C₆alkoxy-C₁-C₆alkyloxy, R₃ is C₁-C₆alkyl, R₄ is C₁-C₆alkyl, and R₅ is C₁-C₆alkyl, C₁-C₆hydroxyalkyl, C₁-C₆alkoxy-C₁-C₆-alkyl, C₁-C₆alkanoyloxy-C₁-C₆alkyl, C₁-C₆aminoalkyl, C₁-C₆alkylamino-C₁-C₆-alkyl, C₁-C₆-dialkylamino-C₁-C₆-alkyl, C₁-C₆-alkanoyl-
15 amido-C₁-C₆-alkyl, HO(O)C-C₁-C₆-alkyl, C₁-C₆alkyl-O-(O)C-C₁-C₆alkyl, H₂N-C(O)-C₁-C₆alkyl, C₁-C₆alkyl-HN-C(O)-C₁-C₆alkyl or (C₁-C₆alkyl)₂N-C(O)-C₁-C₆-alkyl, comprising the steps
a) reaction of a compound of formula II,



20

(II),

with an amine of formula R₅-NH₂ to form a compound of formula III,




(III),

and

b) reduction of the azide group of the compound of formula
 5 III to the amine group and isolation of the compounds of
 formula I, if necessary with the addition of a salt-forming
 acid, comprising the preparation of the compound of formula
 II by reacting

c) a compound of formula IV


10



(IV),

wherein X is Cl, Br or I, with an amine to form a
 carboxamide of formula V,


15



(V),


wherein R₆ is an amino group,

d1) azidating a compound of formula V to form a compound of
 20 formula VI




(VI), or

d2) protecting the hydroxyl group in the compounds of
 formula V, and azidating the resulting compound of formula
 5 VII



(VII),

wherein S is a protecting group, to form a compound of
 10 formula VIII,



(VIII),

e) and then lactonizing the compound of formula VI or VIII
 15 in the presence of an acid to form a compound of formula II.

2. A process according to claim 1 comprising an embodiment
 wherein R₁ is C₁-C₄alkoxy or C₁-C₄alkoxy-C₁-C₄alkyloxy, R₂ is
 C₁-C₄alkoxy, R₃ is C₁-C₄alkyl, R₄ is C₁-C₄alkyl and R₅ is
 20 H₂NC(O)-C₁-C₆alkyl which if necessary is N-monosubstituted or
 N-di-C₁-C₄alkyl substituted.

3. A process according to claim 2 comprising an embodiment
 wherein R₁ is 1-methoxyprop-3-yloxy and R₂ is methoxy.


4. A process according to claim 2 comprising an embodiment wherein R_3 and R_4 are in each case isopropyl.

5. A process according to claim 2 comprising an embodiment wherein R_5 is $H_2NC(O)-C_1-C_6alkyl$.

6. A process according to claim 1 comprising an embodiment wherein R_1 is methoxy- $C_2-C_4alkyloxy$, R_2 is methoxy or ethoxy, R_3 is C_2-C_4alkyl , R_4 is C_2-C_4alkyl and R_5 is $H_2NC(O)-C_1-C_6alkyl$.

7. A process according to claim 1 comprising an embodiment wherein R_1 is 3-methoxy-prop-3-yloxy, R_2 is methoxy, R_3 and R_4 are each 1-methyleth-1-yl, and R_5 is $H_2NC(O)-[C(CH_3)_2]-CH_2-$.

8. Compounds of formula IX,




(IX),

wherein

X is halogen, R_1 and R_2 are, independently of one another, H , C_1-C_6alkyl , $C_1-C_6halogenalkyl$, $C_1-C_6alkoxy$, $C_1-C_6alkoxy-C_1-C_6alkyl$, or $C_1-C_6alkoxy-C_1-C_6alkyloxy$, R_3 is C_1-C_6alkyl , R_4 is C_1-C_6alkyl , R_7 is an amino group, and R_9 is a protecting group or hydrogen.

9. Compounds of formula IXa,



(IXa),

wherein

R₁ and R₂ are, independently of one another, H, C₁-C₆alkyl,
5 C₁-C₆halogenalkyl, C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, or C₁-
C₆alkoxy-C₁-C₆alkyloxy, R₃ is C₁-C₆alkyl, R₄ is C₁-C₆alkyl, R₇
is an amino group, and R₉ is a protecting group or hydrogen.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CH 01/00400

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 C07C231/02 C07C231/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, WPI Data, PAJ, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 678 503 A (CIBA GEIGY AG) 25 October 1995 (1995-10-25) cited in the application page 14, line 1 - line 10 page 23	1,8-12
A	----- PETER HEROLD ET AL: "A Versatile and Stereocontrolled Synthesis of Hydroxyethylene Dipeptide Isomers" JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY, EASTON, US, vol. 54, no. 5, 3 March 1989 (1989-03-03), pages 1178-1185, XP002149098 ISSN: 0022-3263 scheme I page 1179 ----- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

16 October 2001

Date of mailing of the international search report

02/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Bader, K

INTERNATIONAL SEARCH REPORT

International Application No

PCT 01/00400

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 678 500 A (CIBA GEIGY AG) 25 October 1995 (1995-10-25) page 10, line 49 -page 13, line 48 ---	1
A	EP 0 678 514 A (CIBA GEIGY AG) 25 October 1995 (1995-10-25) page 2, line 1 - line 10 page 5, line 18 - line 34 page 23 -page 24; claims 8,9 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CH 01/00400

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0678503	A	25-10-1995	AT 183997 T 15-09-1999
		AU 1642095 A 26-10-1995	
		AU 699616 B2 10-12-1998	
		AU 1642195 A 26-10-1995	
		AU 1642395 A 26-10-1995	
		BR 1100656 A3 06-06-2000	
		CA 2147044 A1 19-10-1995	
		CA 2147052 A1 19-10-1995	
		CA 2147056 A1 19-10-1995	
		CN 1117960 A 06-03-1996	
		CZ 9500976 A3 15-11-1995	
		DE 59506707 D1 07-10-1999	
		DK 678503 T3 20-03-2000	
		EP 0678503 A1 25-10-1995	
		EP 0678514 A1 25-10-1995	
		EP 0678500 A1 25-10-1995	
		ES 2137478 T3 16-12-1999	
		FI 951771 A 19-10-1995	
		FI 951772 A 19-10-1995	
		FI 951773 A 19-10-1995	
		GR 3031997 T3 31-03-2000	
		HU 74074 A2 28-10-1996	
		HU 72110 A2 28-03-1996	
		HU 71701 A2 29-01-1996	
		JP 8053434 A 27-02-1996	
		JP 8081430 A 26-03-1996	
		JP 8027079 A 30-01-1996	
		NO 951441 A 19-10-1995	
		NO 951442 A 19-10-1995	
		NO 951443 A 19-10-1995	
		NZ 270936 A 24-06-1997	
		NZ 270938 A 26-11-1996	
		NZ 270939 A 20-12-1996	
		TW 402582 B 21-08-2000	
		US 5606078 A 25-02-1997	
		US 5659065 A 19-08-1997	
		US 5559111 A 24-09-1996	
		US 5654445 A 05-08-1997	
		US 5646143 A 08-07-1997	
		US 5627182 A 06-05-1997	
		US 5705658 A 06-01-1998	
		ZA 9503050 A 08-11-1995	
		ZA 9503051 A 18-10-1995	
		ZA 9503052 A 18-10-1995	
EP 0678500	A	25-10-1995	AT 183997 T 15-09-1999
		AU 1642095 A 26-10-1995	
		AU 699616 B2 10-12-1998	
		AU 1642195 A 26-10-1995	
		AU 1642395 A 26-10-1995	
		BR 1100656 A3 06-06-2000	
		CA 2147044 A1 19-10-1995	
		CA 2147052 A1 19-10-1995	
		CA 2147056 A1 19-10-1995	
		CN 1117960 A 06-03-1996	
		CZ 9500976 A3 15-11-1995	
		DE 59506707 D1 07-10-1999	
		DK 678503 T3 20-03-2000	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC 01/00400

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0678500	A	EP 0678503 A1	25-10-1995
		EP 0678514 A1	25-10-1995
		EP 0678500 A1	25-10-1995
		ES 2137478 T3	16-12-1999
		FI 951771 A	19-10-1995
		FI 951772 A	19-10-1995
		FI 951773 A	19-10-1995
		GR 3031997 T3	31-03-2000
		HU 74074 A2	28-10-1996
		HU 72110 A2	28-03-1996
		HU 71701 A2	29-01-1996
		JP 8053434 A	27-02-1996
		JP 8081430 A	26-03-1996
		JP 8027079 A	30-01-1996
		NO 951441 A	19-10-1995
		NO 951442 A	19-10-1995
		NO 951443 A	19-10-1995
		NZ 270936 A	24-06-1997
		NZ 270938 A	26-11-1996
		NZ 270939 A	20-12-1996
		TW 402582 B	21-08-2000
		US 5606078 A	25-02-1997
		US 5659065 A	19-08-1997
		US 5559111 A	24-09-1996
		US 5654445 A	05-08-1997
		US 5646143 A	08-07-1997
		US 5627182 A	06-05-1997
		US 5705658 A	06-01-1998
		ZA 9503050 A	08-11-1995
		ZA 9503051 A	18-10-1995
		ZA 9503052 A	18-10-1995
EP 0678514	A	25-10-1995	AT 183997 T
			AU 1642095 A
			AU 699616 B2
			AU 1642195 A
			AU 1642395 A
			BR 1100656 A3
			CA 2147044 A1
			CA 2147052 A1
			CA 2147056 A1
			CN 1117960 A
			CZ 9500976 A3
			DE 59506707 D1
			DK 678503 T3
			EP 0678503 A1
			EP 0678514 A1
			EP 0678500 A1
			ES 2137478 T3
			FI 951771 A
			FI 951772 A
			FI 951773 A
			GR 3031997 T3
			HU 74074 A2
			HU 72110 A2
			HU 71701 A2
			JP 8053434 A
			JP 8081430 A

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CH 01/00400

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0678514	A	JP 8027079	A 30-01-1996
		NO 951441	A 19-10-1995
		NO 951442	A 19-10-1995
		NO 951443	A 19-10-1995
		NZ 270936	A 24-06-1997
		NZ 270938	A 26-11-1996
		NZ 270939	A 20-12-1996
		TW 402582	B 21-08-2000
		US 5606078	A 25-02-1997
		US 5659065	A 19-08-1997
		US 5559111	A 24-09-1996
		US 5654445	A 05-08-1997
		US 5646143	A 08-07-1997
		US 5627182	A 06-05-1997
		US 5705658	A 06-01-1998
		ZA 9503050	A 08-11-1995
		ZA 9503051	A 18-10-1995
		ZA 9503052	A 18-10-1995

THIS PAGE BLANK (USPTO)